



EnterMedSchool.org

Meiosis

Study Guide — Cell Division

Challenging Pre-med style questions on meiosis, ploidy, chromatids/DNA molecules, crossing over, segregation, and independent assortment

24 items — Study Guide with Answers

Free & Open-Source

Licensed under Creative Commons — Attribution required when sharing

Generated February 20, 2026

Scan to visit online





1 Which event occurs in meiosis I but NOT in mitosis of a diploid eukaryotic cell?



- A Separation of sister chromatids at anaphase
- B **Pairing (synapsis) of homologous chromosomes and crossing over in prophase I** ✓
- C Replication of DNA before division
- D Formation of a spindle made of microtubules
- E Alignment of chromosomes at the cell equator

► **Explanation:** Synapsis and crossing over between homologous chromosomes occur only in prophase I of meiosis, not in mitosis.

2 Crossing over in meiosis typically occurs:



- A Between sister chromatids of the same chromosome during prophase I
- B **Between non-sister chromatids of homologous chromosomes during prophase I** ✓
- C Between non-homologous chromosomes during metaphase II
- D Between maternal and paternal chromosomes in G1
- E Between duplicated sister chromatids during cytokinesis

► **Explanation:** A crossover is an exchange of DNA between non-sister chromatids in a homologous pair (a tetrad) during prophase I, creating recombinant chromatids.

3 A diploid species has $2n = 4$. In a primary spermatocyte at metaphase I of meiosis, how many chromosomes and chromatids are present in the cell?



- A 2 chromosomes and 2 chromatids
- B 2 chromosomes and 4 chromatids





- C 4 chromosomes and 4 chromatids
- D 4 chromosomes and 8 chromatids ✓
- E 8 chromosomes and 8 chromatids

► **Explanation:** After S phase, each of the 4 chromosomes has two sister chromatids ($4 \times 2 = 8$ chromatids). Meiosis I does not change chromosome number until after division.

4 The same species ($2n = 4$) completes meiosis I. Immediately after cytokinesis at the end of meiosis I, how many chromosomes and chromatids are present in EACH daughter cell?



- A 2 chromosomes and 2 chromatids
- B 2 chromosomes and 4 chromatids ✓
- C 4 chromosomes and 4 chromatids
- D 4 chromosomes and 8 chromatids
- E 1 chromosome and 2 chromatids

► **Explanation:** Meiosis I separates homologous chromosomes. Each daughter cell is haploid ($n = 2$), but each chromosome is still composed of two sister chromatids ($2 \times 2 = 4$ chromatids).

5 A diploid species has $2n = 6$. In a secondary oocyte at metaphase II, how many chromosomes and chromatids are present in that cell?



- A 3 chromosomes and 3 chromatids
- B 3 chromosomes and 6 chromatids ✓
- C 6 chromosomes and 6 chromatids
- D 6 chromosomes and 12 chromatids
- E 12 chromosomes and 12 chromatids





► **Explanation:** After meiosis I, cells are haploid ($n = 3$), but each chromosome still has two sister chromatids ($3 \times 2 = 6$ chromatids). This is the situation at metaphase II.

6 At which point in meiosis is a cell first considered haploid (n), rather than diploid ($2n$)?



- A Immediately after DNA replication in the pre-meiotic S phase
- B During prophase I when homologous chromosomes pair
- C During metaphase I when homologous pairs line up at the equator
- D After completion of meiosis I, when homologous chromosomes are in different nuclei ✓**
- E Only after completion of meiosis II

► **Explanation:** A nucleus is first haploid when homologous chromosomes have been separated into different nuclei at the end of meiosis I, even though chromosomes are still duplicated.

7 Which statement best distinguishes meiosis I from meiosis II?



- A Meiosis I separates sister chromatids, whereas meiosis II separates homologous chromosomes
- B Meiosis I is reductional, halving the chromosome set by separating homologous chromosomes; meiosis II is equational, separating sister chromatids ✓**
- C DNA replication occurs in both meiosis I and meiosis II
- D Crossing over occurs mainly in meiosis II
- E Meiosis II doubles the chromosome number, whereas meiosis I halves it

► **Explanation:** Meiosis I is reductional ($2n \rightarrow n$) because homologues separate; meiosis II is like mitosis ($n \rightarrow n$), separating sister chromatids.





8 For an autosomal gene with alleles **A** and **a** in a heterozygous individual (**Aa**), which meiotic event most directly explains Mendel's law of segregation?

- A** Replication of the DNA during S phase
- B** Pairing of homologous chromosomes in prophase I
- C** Separation of homologous chromosomes carrying **A** and **a** at anaphase I ✓
- D** Separation of sister chromatids at anaphase II
- E** Random fusion of gametes at fertilisation

► **Explanation:** Alleles **A** and **a** reside on homologous chromosomes; at anaphase I, those homologues separate so each gamete receives only one allele.



9 Which event in meiosis is the physical basis of Mendel's law of independent assortment for genes located on different chromosome pairs?

- A** Random fusion of sperm and egg at fertilisation
- B** Random orientation of each pair of homologous chromosomes on the metaphase I plate ✓
- C** Random positions of crossing-over events along a chromosome in prophase I
- D** Random segregation of sister chromatids at anaphase II
- E** Random occurrence of mutations during DNA replication

► **Explanation:** Each homologous pair can orient in two ways at metaphase I independently of other pairs; this independent alignment is the basis of independent assortment for genes on different chromosomes.



10 A diploid species has $2n = 8$ (four pairs of homologous chromosomes). Ignoring crossing over, how many genetically distinct gamete types can be produced by independent assortment of maternal and paternal chromosomes?





- A 4
- B 8
- C 16 ✓
- D 32
- E 64

► **Explanation:** For n pairs, independent assortment gives 2^n combinations. Here $n = 4$, so $2^4 = 16$ possible combinations of whole maternal/paternal chromosomes.

11 A diploid individual has genotype **AaBb**, where genes **A** and **B** are on different chromosome pairs (unlinked). Assuming normal meiosis, which gamete genotypes and ratios are expected?



- A Only AB and ab, in a 1:1 ratio
- B Only Ab and aB, in a 1:1 ratio
- C **AB, Ab, aB and ab, each with equal probability (1:1:1:1)** ✓
- D AB with 1/2 probability, and ab, Ab, aB each with 1/6 probability
- E A and a only, because B and b segregate in somatic cells

► **Explanation:** For two unlinked heterozygous genes, independent assortment produces four equally frequent gamete types: AB, Ab, aB, ab.

12 An individual has genotype **AB/ab** for two genes that lie close together on the **SAME** chromosome (cis arrangement), and we assume **NO** crossing over occurs between them. Which gametes can be produced with respect to these genes, and in what ratio?



- A Only AB gametes
- B Only Ab gametes





- C AB and ab gametes in a 1:1 ratio ✓
- D AB, Ab, aB and ab gametes in a 1:1:1:1 ratio
- E Ab and aB gametes in a 1:1 ratio

► **Explanation:** If genes are perfectly linked and there is no crossing over between them, each chromatid carries either AB or ab; segregation in meiosis produces only parental combinations AB and ab in equal numbers.

13 Consider an individual with genotype AB/ab for two linked genes on one pair of homologous chromosomes. Suppose that in EVERY meiosis there is exactly ONE crossover between the A and B loci in each tetrad, and all chromatids are equally likely to be included in gametes. What proportion of gametes will carry recombinant haplotypes (Ab or aB)?



- A 0%
- B 25%
- C 50% ✓
- D 75%
- E 100%

► **Explanation:** A single crossover between A and B involves two of the four chromatids: those two become recombinant (Ab and aB), while the other two remain parental (AB and ab). Thus $2/4 = 50\%$ of chromatids, and therefore of gametes, are recombinant.

14 Which statement correctly describes SOURCES of genetic variation in sexually reproducing organisms?



- A Crossing over occurs during fertilisation, while independent assortment occurs in mitosis
- B Independent assortment and crossing over both occur during meiosis; random fertilisation then combines gametes to produce additional variation ✓





- C Independent assortment occurs only when chromosomes fail to pair in meiosis I
- D Crossing over reduces genetic variation by making chromatids more similar
- E Random fertilisation has no effect on genetic variation if independent assortment already occurred

► **Explanation:** Crossing over (recombination) and independent assortment both reshuffle alleles during meiosis; random fertilisation then combines gametes from different parents, further increasing variation.

15 In a diploid organism with $2n = 10$, a germ-line cell has completed DNA replication and is at metaphase I of meiosis. How many DNA molecules (chromatids) are present in this cell?



- A 5
- B 10
- C 20 ✓
- D 40
- E 80

► **Explanation:** $2n = 10$ means 10 chromosomes. After replication each chromosome consists of two sister chromatids, so there are $10 \times 2 = 20$ chromatids (20 long DNA molecules).

16 In the same species ($2n = 10$), how many chromosomes and DNA molecules (chromatids) are present in EACH gamete immediately after meiosis II?



- A 5 chromosomes and 10 DNA molecules
- B 5 chromosomes and 5 DNA molecules ✓
- C 10 chromosomes and 10 DNA molecules
- D 10 chromosomes and 5 DNA molecules





- E 5 chromosomes and 20 DNA molecules

► **Explanation:** Gametes are haploid ($n = 5$) and each chromosome is a single chromatid, so there are 5 chromosomes and 5 DNA molecules per gamete.

17 In a diploid organism, metaphase of mitosis shows $2n$ chromosomes aligned at the equator, each composed of two sister chromatids ($4n$ chromatids total). Which stage of meiosis has the same combination of chromosome and chromatid numbers in a single cell at an alignment stage?



- A Prophase I
- B Metaphase I ✓**
- C Metaphase II
- D Anaphase II
- E Telophase II

► **Explanation:** At metaphase I, the cell is still diploid with $2n$ chromosomes, each consisting of two chromatids ($4n$ chromatids total), now arranged as bivalents on the metaphase I plate.

18 Nondisjunction of ONE pair of homologous chromosomes occurs during meiosis I, but meiosis II proceeds normally. For that chromosome pair, which combination of gametes is produced?



- A Four normal gametes
- B Three normal gametes and one gamete with an extra chromosome ($n + 1$)
- C Two gametes with $n + 1$ and two gametes with $n - 1$ for that chromosome ✓**
- D One gamete with $n + 1$, one with $n - 1$, and two normal gametes
- E Four gametes with $n + 1$





► **Explanation:** If homologues fail to separate in meiosis I, both go to one pole and none to the other. After meiosis II, this yields two gametes with an extra copy ($n + 1$) and two lacking that chromosome ($n - 1$).

19 Nondisjunction of ONE pair of sister chromatids occurs during meiosis II, while meiosis I is normal. For that chromosome pair, which combination of gametes is produced?



- A Four normal gametes
- B Two gametes with $n + 1$ and two with $n - 1$ for that chromosome
- C One gamete with $n + 1$, one with $n - 1$, and two normal gametes ✓**
- D Three normal gametes and one with $n - 1$
- E Four gametes with $n + 1$

► **Explanation:** After a normal meiosis I, each cell has one homologue. Nondisjunction in meiosis II of one cell sends both sister chromatids to one gamete ($n + 1$) and none to the other ($n - 1$), while the other meiosis II division produces two normal gametes.

20 Which statement best describes the effect of crossing over on linked genes (genes on the same chromosome)?



- A Crossing over has no effect on linked genes because they always segregate together
- B Crossing over tends to create new combinations of alleles on a chromosome and can reduce the strength of linkage between genes ✓**
- C Crossing over only occurs between non-homologous chromosomes and therefore cannot affect linked genes
- D Crossing over always guarantees that genes will assort independently, regardless of their distance apart
- E Crossing over doubles the number of chromosomes in a gamete





► **Explanation:** Crossing over between linked genes swaps segments between homologous chromatids, generating recombinant allele combinations and weakening the correlation between genes that would otherwise always co-segregate.

21 During prophase I, homologous chromosomes undergo several substages. In which substage are homologous chromosomes fully synapsed and crossing over actively occurs?



- A Leptotene
- B Zygotene
- C Pachytene ✓**
- D Diplotene
- E Diakinesis

► **Explanation:** In pachytene, the synaptonemal complex is fully formed and crossing over occurs between non-sister chromatids of homologous chromosomes.

22 Which description best defines a chiasma (plural: chiasmata) seen in prophase I and metaphase I?



- A The point where sister chromatids are joined at the centromere
- B The attachment site of spindle microtubules on a chromosome
- C The visible region where non-sister chromatids of homologous chromosomes have exchanged segments due to crossing over ✓**
- D The constriction site where cytokinesis begins
- E The region of the chromosome that always remains heterochromatic

► **Explanation:** A chiasma is the cytological manifestation of a crossover—an X-shaped region where non-sister chromatids have exchanged DNA and still remain temporarily linked.





23 In a species with $2n = 6$, a primary spermatocyte enters metaphase I with three bivalents (tetrads). After a complete meiosis (I and II), which combination correctly describes chromosome and chromatid numbers in (1) that primary spermatocyte at metaphase I, and (2) one of the four resulting sperm cells?



- A (1) 6 chromosomes, 6 chromatids; (2) 3 chromosomes, 6 chromatids
- B (1) 6 chromosomes, 12 chromatids; (2) 3 chromosomes, 3 chromatids ✓**
- C (1) 3 chromosomes, 6 chromatids; (2) 3 chromosomes, 3 chromatids
- D (1) 6 chromosomes, 12 chromatids; (2) 6 chromosomes, 3 chromatids
- E (1) 3 chromosomes, 3 chromatids; (2) 6 chromosomes, 6 chromatids

► **Explanation:** Metaphase I: $2n = 6$ chromosomes, each with two chromatids ($6 \times 2 = 12$). Each sperm cell (haploid, after meiosis II) has $n = 3$ chromosomes, each as a single chromatid ($3 \times 1 = 3$).

24 Which statement about Mendel's laws and chromosome behaviour is correct?



- A The law of segregation depends on crossing over between homologous chromosomes
- B The law of independent assortment applies strictly to all gene pairs, regardless of their positions
- C The law of segregation reflects separation of homologous chromosomes, and the law of independent assortment applies cleanly only to genes on different chromosomes or far apart on the same chromosome ✓**
- D Both laws depend entirely on random mutation events in meiosis
- E The law of segregation is only valid in mitotically dividing cells

► **Explanation:** Segregation is explained by homologues separating in meiosis I. Independent assortment holds exactly for genes on different chromosomes; closely linked genes may violate it unless crossing over separates them often enough.

